



## Emergency Nursing Resource: Gastric Tube Placement Verification

In patients having gastric tubes inserted in the emergency department setting, which bedside technique is best for confirmation of accurate placement immediately after tube insertion compared to radiograph?

### Developed by:

#### 2010 ENA Emergency Nursing Resources Development Committee

Jean A. Proehl, MN, RN, CEN, CPEN, FAEN

Karen Heaton, PhD, FNP-BC, CEN

Mary Kathryn Naccarato, MSN, RN, CEN, HCRM

Melanie A. Crowley, MSN, RN, CEN

Andrew Storer, DNP, RN, CRNP

Jason D. Moretz, BSN, RN, CEN, CTRN

Suling Li, PhD, RN

ENA's Emergency Nursing Resources (ENRs) are developed by ENA members to provide emergency nurses with evidence-based information to utilize and implement in their care of emergency patients and families. Each ENR focuses on a clinical or practice-based issue, and is the result of a review and analysis of current information believed to be reliable. As such, information and recommendations within a particular ENR reflect the current scientific and clinical knowledge at the time of publication, are only current as of their publication date, and are subject to change without notice as advances emerge.

In addition, variations in practice, which take into account the needs of the individual patient and the resources and limitations unique to the institution, may warrant approaches, treatments and/or procedures that differ from the recommendations outlined in the ENRs. Therefore, these recommendations should not be construed as dictating an exclusive course of management, treatment or care, nor does the use of such recommendations guarantee a particular outcome. ENRs are never intended to replace a practitioner's best medical judgment based on the clinical circumstances of a particular patient or patient population. ENRs are published by ENA for educational and informational purposes only, and ENA does not approve or endorse any specific methods, practices, or sources of information. ENA assumes no liability for any injury and/or damage to persons or property arising out of or related to the use of or reliance on any ENR.

**Publication Date:** December 2010

## Table of Contents

|   | Page |
|---|------|
| Background/Significance .....   | 1    |
| Methodology .....   | 1    |
| Evidence Table.....   | 2    |
| Other Resources Table .....   | 2    |
| Summary of Literature Review.....   | 3    |
| Description of Decision Options/Interventions and the Level of Recommendation ..... | 6    |
| Bibliography.....   | 7    |
| Acknowledgement.....  | 9    |

## Background/Significance

Gastric tube (GT) placement is a common bedside procedure performed by registered nurses in the emergency department (ED). Although often considered an innocuous procedure, gastric tube placement can result in serious and even lethal complications such as, misplacement of the gastric tube into the pulmonary system resulting in respiratory distress or death. The standard of care requires verification of the placement of the gastric tube prior to its use in order to minimize complications. Radiographic verification is considered the preferred method of confirmation (Leschke, 2004) and is considered the “gold standard” by many, especially for feeding tubes (Araujo-Preza, Melhado, Gutierrez, Maniatis, & Castellano, 2002; Ellet, 2004; Elpern, Killeen, Talla, Perez, & Gurka, 2007; Kearns & Donna, 2001; Metheny, 2007). However, bedside methods are commonly used as a proxy for radiographic verification when large bore GTs are inserted because of the associated cost, time delay, and radiation exposure. In addition, a radiographic test cannot be performed by the bedside nurse. It has been well documented for almost 20 years that a common bedside method (auscultation) is often inaccurate; however, it is still widely practiced. This Emergency Nursing Resource (ENR) aims to evaluate various bedside gastric tube placement verification methods as an alternative to radiography.

## Methodology

This ENR was created based on a thorough review and critical analysis of the literature following ENA’s [Guidelines for the Development of the Emergency Nursing Resources](#). Via a comprehensive literature search, all articles relevant to the topic were identified. The following databases were searched: PubMed, eTBLAST, CINAHL, Cochrane - British Medical Journal, Agency for Healthcare Research and Quality (AHRQ; [www.ahrq.gov](http://www.ahrq.gov)), and the National Guideline Clearinghouse ([www.guidelines.gov](http://www.guidelines.gov)). Searches were conducted using the key words nasoenteral tubes, tube placement determination, gastric tubes, gastric tube placement confirmation, gastric tube placement, and nasoenteral tubes + catheters and tubes. Initial searches were limited to English language articles on human subjects from 2005 – October, 2010. This five year search limit was found to be inadequate so the time frame was expanded to 1994 and a specific search was performed for Metheny’s publications because of the seminal nature of her work. In addition, the reference lists in the selected articles were scanned for pertinent research findings. Research articles from emergency department settings, non-ED settings, position statements and guidelines from other sources were also reviewed.

Articles that met the following criteria were chosen to formulate the ENR: research studies, meta-analyses, systematic reviews, and existing guidelines relevant to the topic. Other types of article were also reviewed and provided as additional information. The ENR authors used [standardized worksheets](#), including Evidence-Appraisal Table Template, Critique Worksheet and AGREE Work Sheet, to prepare tables of evidence ranking each article in terms of the level of evidence, quality of evidence, and relevance and applicability to practice. Clinical findings and levels of recommendations regarding patient management were then made by the Emergency Nursing Resource Development Committee according to the ENA’s classification of levels of recommendation for practice, which include: Level A High, Level B. Moderate, Level C. Weak or Not recommended for practice (See Table 1).

**Table 1. Levels of Recommendation for Practice**

|   |
|---|
| <p><b><u>Level A recommendations: High</u></b></p> <ul style="list-style-type: none"> <li>• Reflects a high degree of clinical certainty</li> <li>• Based on availability of high quality level I, II and/or III evidence available using Melnyk &amp; Fineout-Overholt grading system (Melnyk &amp; Fineout-Overholt, 2005)</li> <li>• Based on consistent and good quality evidence; has relevance and applicability to emergency nursing practice</li> <li>• Is beneficial</li> </ul>  |
| <p><b><u>Level B recommendations: Moderate</u></b></p> <ul style="list-style-type: none"> <li>• Reflects moderate clinical certainty</li> <li>• Based on availability of Level III and/or Level IV and V evidence using Melnyk &amp; Fineout-Overholt grading system (Melnyk &amp; Fineout-Overholt, 2005)</li> <li>• There are some minor or inconsistencies in quality evidence; has relevance and applicability to emergency nursing practice</li> <li>• Is likely to be beneficial.</li> </ul>  |
| <p><b><u>Level C recommendations: Weak</u></b></p> <ul style="list-style-type: none"> <li>• Level V, VI and/or VII evidence available using Melnyk &amp; Fineout-Overholt grading system (Melnyk &amp; Fineout-Overholt, 2005) - Based on consensus, usual practice, evidence, case series for studies of treatment or screening, anecdotal evidence and/or opinion</li> <li>• There is limited or low quality patient-oriented evidence; has relevance and applicability to emergency nursing practice</li> <li>• Has limited or unknown effectiveness</li> </ul>  |
| <p><b><u>Not recommended for practice</u></b></p> <ul style="list-style-type: none"> <li>• No objective evidence or only anecdotal evidence available; or the supportive evidence is from poorly controlled or uncontrolled studies</li> <li>• Other indications for not recommending evidence for practice may include: <ul style="list-style-type: none"> <li>○ Conflicting evidence</li> <li>○ Harmfulness has been demonstrated</li> <li>○ Cost or burden necessary for intervention exceeds anticipated benefit</li> <li>○ Does not have relevance or applicability to emergency nursing practice</li> </ul> </li> <li>• There are certain circumstances in which the recommendations stemming from a body of evidence should not be rated as highly as the individual studies on which they are based. For example: <ul style="list-style-type: none"> <li>○ Heterogeneity of results</li> <li>○ Uncertainty about effect magnitude and consequence</li> <li>○ Strength of prior beliefs</li> <li>○ Publication bias</li> </ul> </li> </ul> |

**Evidence Table and Other Resources**

The articles reviewed to formulate the ENR are described in the [Evidence Table](#). Other articles relevant to GT placement were reviewed to serve as additional resources ([Other Resources Table](#)).

## **Summary of Literature Review**

### **Gastric Tubes**

Gastric tubes may be inserted nasally, i.e. nasogastric tubes, or orally, i.e. orogastric tubes. Regardless of the insertion route all are GTs so this term will be used throughout the document. The main reasons for inserting a GT in the emergency department are to decompress the stomach and remove stomach contents; to prevent aspiration and minimize nausea/vomiting; or to instill liquids or medications (Christensen, 2001). Two categories, large and small bore gastric tubes have been designed to meet these treatment needs. For example, large bore GTs are considered for short term use and aid in the removal of stomach contents and the instillation of liquids or medications. In contrast, small bore GTs, also known as feeding tubes, remain in place for a longer period of time and are reserved for the instillation of enteral nutrition, liquids and medications. Thus, large bore GTs, not feeding tubes, are typically inserted in the emergency department.

Treatment needs guide the decision about the location of the tip of the gastric tube. A large bore gastric tube is inserted via the nose or mouth and guided into the stomach. Whereas, a small bore gastric tube is advanced through the stomach into the small intestine. Because anatomical changes associated with growth and development occur; patient age and size are also considered when determining the depth of insertion and size of the gastric tube selected (Cincinnati Guidelines, 2009).

Although most studies of gastric tube bedside verification methods focus on small bore feeding tubes; the limited numbers of studies conducted in emergency department settings using large bore gastric tubes are also included in this review. The most common bedside verification methods can be categorized as non-aspirate or aspirate.

### **Non-Aspirate Methods**

Non-aspirate methods include auscultation, carbon dioxide detection, transillumination, and magnetic detection; whereas, aspirate methods are: visual characteristics, pH, bilirubin, and enzyme tests.

#### ***Auscultation***

The auscultatory method involves instillation of air into the tube while simultaneously listening with a stethoscope for a sound of air over the epigastric region. Auscultation alone continues to be used by nurses currently caring for neonates (Cincinnati Guidelines, 2009), pediatric patients (de Boer, 2009) and adults (Nursing, 2006) despite its proven unreliability as a single verification method (Ellett, 2005; Metheny, 1994; Neuman, 1995; Metheny, 1999; Yardley & Donaldson, 2010). Results from a 2006 on-line survey of 1,600 nurses indicated that 65% used the auscultation verification method most of the time (Nursing). Guidelines published by Cincinnati Children's Hospital (2009) reported only 60-80% reliability with auscultation as a single verification method. An American Association of Critical Care Nurses (AACN) practice alert in 2007 suggested abandoning the auscultatory method for gastric tube placement verification due to its unreliability. Thus, pursuit of a reliable, valid, bedside verification method for gastric tube placement has led researchers to investigate methods other than auscultation.

### ***Carbon Dioxide Detection Monitoring***

Misplacement of the gastric tube into the pulmonary system warrants immediate and accurate detection. Studies using CO<sub>2</sub> detection methods (CO<sub>2</sub> monitoring/capnography) were conducted to identify a method that detects gastric tube misplacement (Burns et al., 2006; Elpern et al., 2007). Burns and colleagues (2006) reported 100% agreement between colorimetry and capnography in the identification of CO<sub>2</sub> when the gastric tube was placed inside an endotracheal tube (in situ). Further, 130 adult medical intensive care unit patients underwent large bore GT placement and insertion failure or GT misplacement, was correctly identified by capnography, in 52 patients (a rate of 27%) (Burns, et al. 2006). Gastric tube insertion failures were associated with nasal insertion route ( $p = 0.03$ ) and among spontaneously breathing/non-intubated patients ( $p = 0.01$ ). The small number of misplaced GTs limits the generalizability of the study results.

Capnometry and air insufflations/auscultation were compared to abdominal radiograph for accuracy in detecting misplaced GTs during initial insertion in 91 adult critical care and telemetry patients (Elpern, 2006). Elpern and colleagues reported a 100% success rate in placing the GT into the stomach. However, when compared with abdominal radiographs, 16% of correctly placed GTs via capnography were false positives (indicated to be in the pulmonary track but actually in the GI tract) and there were 5% false positive results with air insufflation/auscultation (Elpern, 2007). While a false positive reading does not immediately jeopardize patient safety, it does require the use of additional verification methods to ensure tube location. Study limitations included sample size, adult-only study population, and false positive readings.

Further research is needed to determine the role of carbon dioxide detection in GT placement. Carbon dioxide detection and monitoring equipment is commonly found in the emergency department because of its use with endotracheal intubation and sedation, however its use with GT placement in the emergency department remains under studied. Interestingly, two GT verification algorithms (Cincinnati, 2009; Metheny, 2001) do not include the carbon dioxide detection method. Instead, both of these algorithms suggest the nurse listen for air movement and/or observe for respiratory distress signs and symptoms to detect the misplacement of the GT.

### ***Transillumination and Magnetic Detection***

Research has also been conducted to determine the feasibility of using transillumination or magnetic detection for GT placement verification. One study utilized a fiberoptic method for GT placement verification (Rulli, 2007). A flexible fiberoptic cable was inserted into the GT of 16 patients, 8 adults and 8 children, who were undergoing a surgical procedure. Transillumination of the epigastric abdominal area was used to indicate correct placement of the GT. Gastric tube placement was confirmed in 100% of the patients. While the study was highly relevant; limitations included small sample size, lack of commercially available equipment, and the operating room practice setting.

Magnetic detection was used to detect position of GT in 88 volunteer subjects, aged 18-75 years (Tobin, 2000). A commercial feeding tube was modified to substitute a magnet for the tungsten weights in the tip of the GT. Prototype magnet detectors determined real-time GT location, orientation and depth of distal end of the feeding tube. Gastric tubes were determined, by fluoroscopy, to be below the diaphragm 100% of the time. A prospective blinded trial of 134 patients compared four GT verification methods: electromagnetic technique, auscultation, aspiration and pH (Kearns & Donna, 2001). Electromagnetic and aspiration method identified tubes above the diaphragm. Electromagnetic method

successfully identified GT placement 90% of the time compared to 53% successful placement using aspiration. Several study limitations included lack of commercially available equipment for both GT and magnetic field detector, laboratory setting, and lack of testing of misplacement of GT in the pulmonary system.

### **Ultrasound**

Ultrasound is clinically feasible because many EDs now have bedside ultrasound units, but research is necessary to validate this method for large-bore GTs in the ED setting. There are limited data emerging regarding the use of ultrasound for confirmation of gastric and feeding tube placement in adults (Nikandros, Skampas, Theodorakopoulou, Ioannidou, Theotokas, & Armaganidis, 2006; Vigneau, Baudel, Guidet, Offenstadt, & Maury, 2005). While this technique looks promising for verification of feeding tube placement, there are no data on the use of this verification method for large bore GTs in the ED setting.

### **Aspirate Methods**

Tests evaluating aspirate content offer an alternative method for verifying GT placement. Visual inspection and biochemical markers such as: pH, bilirubin, and enzymes were the most frequent aspirate methods used to study GT placement. Visual inspection of aspirate involved differentiating appearance of the aspirate obtained from the stomach, small intestine, and lung contents. Metheny and colleagues (1994) reported that critical care nurses were able to differentiate between gastric and intestinal aspirate appearances 90.47% of the time ( $p < 0.01$ ); yet were unable to distinguish between gastric and pulmonary aspirate. Reliable verification methods are needed to determine tube misplacement into the pulmonary system since this is the most common and potentially lethal site for misplacement.

Several studies investigated the biochemical markers of pH, bilirubin, pepsin, and trypsin for GT placement (Cincinnati, 2009; Ellett, 2005; Kearns & Donna 2001; Metheny, 1989, 1994, 1997, 1999; Metheny & Titler, 2001; Phang, Marsh, Barlows, & Schwartz, 2004; Stock, Gilbertson & Babl, 2008; Taylor & Clemente, 2005). Small bore feeding tubes were utilized in all studies. A combination of small bore feeding tubes and large bore GT for decompression were utilized in the Stock, Gilbertson, & Babl, 2008 study and the Cincinnati, Guidelines also address large bore tubes. In addition, study populations often included patients receiving acid inhibiting medications.

Biochemical marker threshold values varied among the studies ranging from a gastric pH value of less than 3.9 to 7 and bilirubin less than 5 to differentiate GT placement in the stomach versus the pulmonary system. Participants in these studies received acid suppressive therapy and tube feedings-both of which influence gastric pH. Reliability of pH testing to determine tube placement within the gastrointestinal tract ranged from 84% (Stock, Gilbertson, & Babl, 2008) to 97% (Metheny, 2000), compared to bilirubin test reliability of 91% (Metheny, 1999), pepsin enzyme reliability of 91.2% and trypsin enzyme reliability of 91.8% (Metheny, Stewart, Smith, & Yan, 1997). Study results reported an alteration in pH test results for patients receiving acid suppression medication. In fact, Taylor and Clemente (2005) reported a 58% reduction in pH confirmation of GT placement for patients receiving acid inhibiting medications.

Research in 2000 by Metheny and colleagues, reported the combined test results of pH greater than 5 and bilirubin less than 5 successfully identified 98.6% of the 141 cases as gastric placement. Laboratory-

based testing of bilirubin, pepsin and trypsin, limit their use as bedside point of care methods. There is limited information using urine bilirubin test strips for the purpose of bedside verification (Metheny, Stewart, Smith, & Yan, 2000) while bedside testing of gastrointestinal enzymes awaits development.

### **Combined Non-Aspirate and Aspirate Methods**

#### ***Algorithmic Approach***

The rate of GT placement accuracy increases when combining non-radiological verification methods rather than reliance on a single bedside verification method (Cincinnati, 2009; Metheny, 2001). Metheny (2001) recommends an algorithm for GT placement verification. Metheny's algorithm for newly inserted large-bore GT begins with auscultation followed by pH and visual inspection of aspirate. The Cincinnati guidelines (2009) also use an algorithm consisting of non-aspirate and aspirate verification methods of auscultation, visual, and pH testing. Study results indicated GT placement achieved a probable accuracy of 97-99% when using auscultation, visual aspirate inspection, and aspirate pH testing (Cincinnati, 2009). There is evidence to support use of a combination of methods of bedside verification for GT placement; however additional research is needed to determine which methods are the most accurate and in what sequence they should be used.

### **Description of Decision Options/Interventions and the Level of Recommendation**

Conclusion and recommendations about initial GT placement bedside verification methods in the emergency department:

- Auscultation as a single verification method is unreliable in determining GT placement (Not recommended)
- There is insufficient evidence to support the use of carbon dioxide detection methods as a single GT placement bedside verification method (Level C: Weak)
- There is insufficient evidence to support the use of transillumination and magnetic detection methods along with equipment and laboratory setting limitations (Level C: Weak)
- There is sufficient evidence to support pH testing of GT aspirates a component of a multiple method bedside verification approach (Level B: Moderate)

## Bibliography

- Cincinnati Children's Hospital Medical Center (CCHMC) (2009). NGT Placement Confirmation, BEST #024. In CCHMC (Ed.), (pp. 1-11).
- de Boer, J., & Smit, B. J. (2009). Nasogastric tube position and intragastric air collection in a neonatal intensive care population. *Advances in Neonatal Care, 9*, 293-298.
- Ellett, M. L. C., Croffie, J. M. B., Cohen, M. D., & Perkins, S. M. (2005). Gastric tube placement in young children. *Clinical Nursing Research, 14*, 238-252.
- Ellett, M. L. C. (2004). What is known about methods of correctly placing gastric tubes in adults and children. [CEU]. *Gastroenterology Nursing, 27*(6), 253-261.
- Elpern, E. H., Killeen, K., Talla, E., Perez, G., & Gurka, D. (2007). Capnometry and air insufflation for assessing initial placement of gastric tubes. *American Journal of Critical Care, 16*(6), 544-550.
- Kearns, P. J., & Donna, C. (2001). A controlled comparison of traditional feeding tube verification methods to a bedside, electromagnetic technique. *Journal of Parenteral and Enteral Nutrition, 25*(4), 210-215.
- Leschke, R. Chapter 47: Nasogastric Intubation in Reichman, E. & Simon, R. (eds.). *Emergency Medicine Procedures*: <http://www.accessemergencymedicine.com/content.aspx?aID=50768>.
- Metheny, N., Reed, L., Berglund, B., & Wehrle, M. A. (1994). Visual characteristics of aspirates from feeding tubes as a method for predicting tube location. *Nursing Research 43*(5), 282-287.
- Metheny, N., & Clouse, R. E. (1997). Bedside methods for detecting aspiration in tube-fed patients. *Chest, 111* (724-731), 724.
- Metheny, N., Stewart, B. J., Smith, L., & Yan, H. (1997). pH and concentrations of pepsin and trypsin in feeding tube aspirates as predictors of tube placement. *Journal of Parenteral and Enteral Nutrition, 21*(5), 279-285
- Metheny, N., Smith, L., & Stewart, B. J. (2000). Development of a reliable and valid bedside test for bilirubin and its utility for improving prediction of feeding tube location. *Nursing Research, 49*(6), 302-309.
- Metheny, N., & Titler, M. G. (2001). Assessing placement of feeding tubes. *American Journal of Nursing, 101*(5), 36-46.
- Metheny, N., Schnelker, R., McGinnis, J., Zimmerman, G., Duke, C., Merritt, B., et al. (2005). Indicators of tubesite during feedings. *Journal of Neuroscience Nursing, 37*(6), 320-326.
- Nursing. (2006). Survey responses: confirming placement of NG tubes. *Nursing 2006, 36*(1), 34.
- Phang, J. S., Marsh, W. A., Barlows, T. G., & Schwartz, H. I. (2001). Determining feeding tube location by gastric and intestinal pH values. *Nutrition in Clinical Practice, 19*, 640-644.
- Rulli, F., Galata, G., Villa, M., Maura, A., Ridolfi, C., Grande, M., et al. (2007). A simple indicator of correct nasogastric suction tube placement in children and adults. *Endoscopy, 39*, E237-E238.
- Stock, A., Gilbertson, H., & Babl, F. E. (2008). Confirming nasogastric tube position in the emergency department: pH testing is reliable. *Pediatric Emergency Care, 24*(12), 805-809
- Taylor, S. J., & Clemente, R. (2005). Confirmation of nasogastric tube position by pH testing. *Journal of Human Nutrition and Dietetics, 18*, 371-375.
- Nikandros, M., Skampas, N., Theodorakopoulou, M., Ioannidou, S., Heotokas, M., & Armaganidis, A. (2006). Sonography as a tool to confirm the position of the nasogastric tube in ICU patients (abstract). *Critical Care, 10* (Suppl 1), S90.
- Tobin, R. W., Gonzales, A. J., Golden, R. N., Brown, M. C., & Silverstein, F. E. (2000). Magnetic detection to position of human nasogastric tubes. *Biomedical Instrumentation and Technology, 34*, 432-436.
- Vigneau, C., Baudel, J., Guidet, B., Offenstadt, G., & Maury, E. (2005). Sonography as an alternative to radiography for nasogastric feeding tube location. *Intensive Care Medicine, 31*, 1570-1572.

Wilkes-Holmes, C. (2006). Safe placement of nasogastric tubes in children. *Paediatric Nursing, 18*(9), 14-17.

Yardley, I. E., & Donaldson, L. J. (2010). Patient safety matters: reducing the risks of nasogastric tubes. *Clinical Medicine, 10*, 228-30.

© Emergency Nurses Association, 2010.

## **Acknowledgement**

ENA would like to acknowledge the 2010 Institute for Emergency Nursing Research (IENR) Advisory Council for the review of this document. Members of the IENR Advisory Council include:

Susan Barnason, PhD, RN, APRN-CNS, CEN, CCRN, FAHA  
Cynthia Dakin, PhD, RN  
Gordon Gillespie, PhD, RN, PHCNS-BC, CEN, CPEN, CCRN, FAEN  
Vicki Keough, PhD, RN, ACNP, CCRN  
Anne Manton, PhD, RN, PMHNP-BC, FAEN, FAAN  
Lisa Wolf, MS, RN, CEN  
Mary Kamienski, PhD, RN, APN, CEN, FAEN, ENA Board of Directors Liaison

ENA also acknowledges Jill Walsh, DNP, RN, ENA Chief Nursing Officer, for the support of this project and the assistance of Leslie Gates in coordinating the work of the Committee.