



# Position Statement

930 E. Woodfield Road, Schaumburg, IL 60173 | 800.900.9659 | ena.org

## Violence and Its Impact on the Emergency Nurse

### Description

In 2002, the World Health Organization (WHO) declared workplace violence (WPV) to be a global epidemic with a negative impact on retention of health personnel and delivery of healthcare.<sup>1</sup> The violence also results in significant economic, personal, and professional costs.<sup>1-3</sup> In the U.S., the prevalence of WPV in the healthcare industry is four times higher than in other private industries.<sup>4</sup> Ease of public access, crowding, long wait times, presence of weapons, and other factors make the emergency department (ED) a highly vulnerable area,<sup>5-9</sup> especially where triage occurs.<sup>10,11</sup> Emergency nurses and other ED staff have a serious occupational risk for WPV, including both verbal and physical assaults.<sup>5-7</sup> For these reasons, WPV has been recognized in many states as a violent crime.<sup>12</sup> Yet, at the time of this publication, only about 30 states have adopted laws making it a felony to assault a registered nurse.<sup>13</sup> Other ongoing legislative initiatives include the introduction of H.R. 1309, Workplace Violence Prevention of Health Care and Social Service Workers Act, in 2018 and continued advocacy and evaluation of state-based felony reforms.

There are no standard definitions of WPV. The Occupational Safety and Health Administration (OSHA) defines WPV as "...any act or threat of physical violence, harassment, intimidation, or other threatening disruptive behavior that occurs at the work site."<sup>39(p1)</sup> WHO uses a broader global definition that encompasses "Incidents where staff are abused, threatened or assaulted in circumstances related to their work, including commuting to and from work, involving an explicit or implicit challenge to their safety, well-being or health".<sup>1p.3</sup> Both definitions demonstrate that WPV manifests in myriad ways as emotional or verbal abuse, coercive or threatening behavior, or physical and sexual assault,<sup>4</sup> and can involve consumers, providers, and organizations.<sup>40</sup> The patient population (e.g., active substance use), along with work schedule (i.e., night shift) experience level, and younger age of the healthcare provider, are consistent risk factors for WPV.<sup>8,10,14-17</sup> Acts of WPV can cause physical and/or psychological harm to emergency nurses leading to job dissatisfaction, emotional exhaustion, burnout, secondary trauma stress, PTSD, absenteeism, and intentions to leave the job or nursing profession,<sup>4,14-24</sup> all of which have potential impacts on patient care due to nurses' decreased productivity, organizational commitment, and engagement.<sup>16,17,24-26</sup> WPV is increasingly seen as a contributing driver of poor nurse retention and recruitment, further exacerbating the nursing shortage and its costly consequences for healthcare organizations and their patients.<sup>4,17,19,24,26-28</sup>

Despite continued education, legislation, and research to increase awareness and understanding of the issue, emergency nurses are reluctant to report incidents of WPV because they believe it is not violence if they did not sustain an injury, reporting can be laborious and futile, patients are not seen as responsible because of their age or illness, and WPV is an expected part of the job.<sup>22,27</sup> Different types of violence exist independently, overlap, and enable each other. For example, relational WPV (bullying) may impede early recognition and management of the violent person because it contributes to nurses' burnout and emotional and physical fatigue.<sup>18,20,27</sup> Similarly, organizations knowingly and unnecessarily exposing their



# Position Statement

930 E. Woodfield Road, Schaumburg, IL 60173 | 800.900.9659 | ena.org

workers to violent situations or allowing a climate of abuse, bullying, or incivility to thrive in the workplace (organizational WPV) can create an environment wherein both relational and consumer violence are ignored, allowing the behaviors to continue without administrative intervention.<sup>10,18,26,28,33</sup>

Researchers suggest an increased emphasis on improving the practice environment (e.g., modifications to physical features to increase visibility or use of patient alert flags) to facilitate adequate assessment, recognition, and shared communication to improve management of the potentially violent person. They also suggest adopting best practices such as increased staff training in stress reduction, conflict resolution, simplified reporting, de-escalation, and behavior management skills, along with a focus on prevention and mitigation of all types of WPV rather than solely on post-incident response to and management of WPV sequelae.<sup>9,16-18,28,29,33-37,41</sup>

## ENA Position

It is the position of the Emergency Nurses Association (ENA) that:

1. Emergency nurses are at significant occupational risk for WPV.
2. The mitigation of WPV requires a zero-tolerance environment instituted and supported by hospital leadership.
3. Emergency nurses have the right to personal safety in the work environment.
4. Emergency nurses have the right to education and training related to the recognition, management, and mitigation of all types of WPV.
5. Emergency nurses have the right and responsibility to report incidents of WPV to their employer and law enforcement without reprisal.
6. Emergency nurses have the right to expectations of privacy, appropriate injury care, and the option for debriefing and professional counseling.
7. Protection against acts of violence include effective administrative, environmental, educational, and security components.
8. Emergency nurses advocate for adoption or continuation of state and federal legislation focused on the prevention of WPV and protection of emergency nurses.
9. Emergency nurses have a vested interest in, and a responsibility to conduct and participate in, research and quality improvement initiatives aimed at preventing, mitigating, and reporting all types and forms of WPV.



# Position Statement

930 E. Woodfield Road, Schaumburg, IL 60173 | 800.900.9659 | ena.org

## Background

To increase program effectiveness, it is recommended that a WPV prevention program include training; formal incident reporting procedures; and administrative, environmental and consumer risk assessment, physical design, and security components to address all types of violence.<sup>3,4,6,27,29,30,34–37,41</sup> When establishing a WPV prevention program, WPV experts recommend that healthcare organizations adopt a multi-faceted, collaborative, interdisciplinary approach that includes a variety of stakeholders such as healthcare administrators, ED managers, clinicians and staff, law enforcement and security personnel, and specialty providers such as mental health practitioners.<sup>27,28,33–35,37</sup> Given the crucial focus on prevention of WPV by patients, visitors, coworkers, and intimate partners, coordination and advocacy among employees, healthcare employers, managers, and nursing leadership is considered necessary for effective implementation of educational, administrative, behavioral, legislative, and engineering approaches necessary for mitigating WPV.<sup>3,4,30,35–37</sup>

Emergency nurses, with their high risk for experiencing WPV, can serve an integral role in all aspects of violence prevention, planning, monitoring, and reporting. Underreporting is a documented barrier to effective identification and mitigation of workplace violence, although nurses do report incidents using both informal (e.g., telling a supervisor or colleague) and formal channels — the latter being more likely when an injury is sustained.<sup>29,30,32</sup> Studies have found that nurses' reluctance to report involves fear of retaliation or dismissal from perpetrators, whether they are patients, colleagues, or supervisors; or community members such as law enforcement<sup>29–31</sup> however, it may be intensified when there is a power imbalance (e.g., a supervisor bullying an employee).<sup>24,26</sup> Organizational commitment to reducing WPV—including workplace policies such as zero tolerance and the role of positive nursing leadership for establishing a non-punitive just culture that discourages bullying and retaliation—is essential to mitigating all types and forms of WPV in emergency nursing and other healthcare environments; a just culture supports the nurse's right to privacy in reporting, injury care, debriefing, and counseling.<sup>18,21,24,26–28</sup>

Further research is essential to determine effective prevention and mitigation strategies, educational priorities for nurse recognition of potential high-risk patients, and conditions for the proactive reduction of WPV. Reporting deficiencies, research design and methodology issues, and inconsistencies in definitions of violence (e.g., threats, assaults, battery) make it difficult to evaluate and compare results across studies both in the United States and globally.<sup>3,7,24,38</sup> Accurate risk surveillance, successful mitigation, and evaluation of WPV interventions will not be possible without standardized definitions, more rigorous data-driven studies, and improved incident reporting, making this an important focus for future WPV research.<sup>3,24,30,37,38,41</sup>

## Resources

American College of Emergency Physicians. (2018). Violence in the emergency department: Resources for a safer workplace. Retrieved from <https://www.acep.org/administration/violence-in-the-emergency-department-resources-for-a-safer-workplace/>



# Position Statement

930 E. Woodfield Road, Schaumburg, IL 60173 | 800.900.9659 | ena.org

Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health. (2018). *Workplace violence prevention for nurses*. CDC Course No. WB2908 – NIOSH Pub. No. 2013-155. [https://www.cdc.gov/niosh/topics/violence/training\\_nurses.html](https://www.cdc.gov/niosh/topics/violence/training_nurses.html)

Emergency Nurses Association. (2019). Workplace violence. <https://www.ena.org/practice-resources/workplace-violence>

U.S. Department of Labor, Office of the Assistant Secretary for Administration & Management. (2019). DOL workplace violence program. Retrieved from <https://www.dol.gov/agencies/oasam/human-resources-center/policies/workplace-violence-program>

## References

1. World Health Organization. (2002). *Framework guidelines for addressing workplace violence in the health sector*. Retrieved from [https://www.who.int/violence\\_injury\\_prevention/violence/interpersonal/en/WVguidelinesEN.pdf?ua=1](https://www.who.int/violence_injury_prevention/violence/interpersonal/en/WVguidelinesEN.pdf?ua=1)
2. Nikathil, S., Olausson, A., Gocentas, R. A., Symons, E., & Mitra, B. (2017). Review article: Workplace violence in the emergency department: A systematic review and meta analysis. *Emergency Medicine Australasia*, 29(3), 265–275. <https://doi.org/10.1111/1742-6723.12761>
3. Phillips, J. P. (2016). Workplace violence against healthcare workers in the United States. *New England Journal of Medicine*, 374(17), 1161–1669. <https://doi.org/10.1056/NEJMra1501998>
4. Occupational Health and Safety Administration. (2015). *Workplace violence in healthcare: Understanding the challenge*. Retrieved from <https://www.osha.gov/Publications/OSHA3826.pdf>
5. Gacki-Smith, J., Juarez, A. M., Boyett, L., Homeyer, C., Robinson, L., & Maclean, S. L. (2009). Violence against nurses working in U.S. emergency departments. *JONA: The Journal of Nursing Administration*, 39(7–8), 340–348. <https://doi.org/10.1097/NNA.0b013e3181ae97db>
6. The Joint Commission. (2018). Physical and verbal violence against health care workers. *Sentinel Event Alert*, 59, 1–9. Retrieved from [https://www.jointcommission.org/assets/1/18/SEA\\_59\\_Workplace\\_violence\\_4\\_13\\_18\\_FINAL.pdf](https://www.jointcommission.org/assets/1/18/SEA_59_Workplace_violence_4_13_18_FINAL.pdf)
7. Spector, P. E., Zhou, Z. E., & Che, X. X. (2014). Nurse exposure to physical and nonphysical violence, bullying, and sexual harassment: A quantitative review. *International Journal of Nursing Studies*, 51(1), 72–84. <https://doi.org/10.1016/j.ijnurstu.2013.01.010>
8. Kleissl-Muir, S., Raymond, A., & Rahman, M. A. (2018). Incidence and factors associated with substance abuse and patient-related violence in the emergency department: A literature review. *Australasian Emergency Care*, 21(4), 159–170. <https://doi.org/10.1016/j.auec.2018.10.004>
9. Stowell, K. R., Hughes, N. P., & Rozel, J. S. (2016). Violence in the emergency department. *Psychiatric Clinics of North America*, 39(4), 557–566. <https://doi.org/10.1016/j.psc.2016.07.003>
10. Pich, J. V., Kable, A., & Hazelton, M. (2017). Antecedents and precipitants of patient-related violence in the emergency department: Results from the Australian VENT study (Violence in Emergency Nursing and Triage). *Australasian Emergency Nursing Journal*, 20(3), 107–113. <https://doi.org/10.1016/j.aenji.2017.05.005>
11. Lau, J. B. C., Magarey, J., & Wiechula, R. (2012). Violence in the emergency department: An ethnographic

# Position Statement

930 E. Woodfield Road, Schaumburg, IL 60173 | 800.900.9659 | ena.org

- study (part II). *International Emergency Nursing*, 20(3), 126–132. <https://doi.org/10.1016/j.ienj.2011.08.001>
12. Papa, A., & Venella, J. (2013). Workplace violence in healthcare: Strategies for advocacy. *The Online Journal of Issues in Nursing*, 18(1), 5. <https://doi.org/10.3912/OJIN.Vol18No01Man05>
  13. Emergency Nurses Association. (2019). States with enhanced penalties for workplace violence against emergency nurses. Retrieved from [https://www.ena.org/docs/default-source/government-relations/ena-wpv-map-2019.pdf?sfvrsn=ad0e8b60\\_8](https://www.ena.org/docs/default-source/government-relations/ena-wpv-map-2019.pdf?sfvrsn=ad0e8b60_8)
  14. Taylor, J. L., & Rew, L. (2011). A systematic review of the literature: Workplace violence in the emergency department. *Journal of Clinical Nursing*, 20(7–8), 1072–1085. <https://doi.org/10.1111/j.1365-2702.2010.03342.x>
  15. Emergency Nurses Association. (2011). *Emergency department violence surveillance study*. Schaumburg, IL: Author. Retrieved from [https://www.ena.org/docs/default-source/resource-library/practice-resources/workplace-violence/2011-emergency-department-violence-surveillance-report.pdf?sfvrsn=5ad81911\\_4](https://www.ena.org/docs/default-source/resource-library/practice-resources/workplace-violence/2011-emergency-department-violence-surveillance-report.pdf?sfvrsn=5ad81911_4)
  16. Stowell, K. R., Hughes, N. P., & Rozel, J. S. (2016). Violence in the emergency department. *Psychiatric Clinics of North America*, 39(4), 557–566. <https://doi.org/10.1016/j.psc.2016.07.003>
  17. Speroni, K. G., Fitch, T., Dawson, E., Dugan, L., & Atherton, M. (2014). Incidence and cost of nurse workplace violence perpetrated by hospital patients or patient visitors. *Journal of Emergency Nursing*, 40(3), 218–228. <https://doi.org/10.1016/j.jen.2013.05.014>
  18. Ashton, R. A., Morris, L., & Smith, I. (2018). A qualitative meta-synthesis of emergency department staff experiences of violence and aggression. *International Emergency Nursing*, 39, 13–19. <https://doi.org/10.1016/j.ienj.2017.12.004>
  19. Kowalenko, T., Gates, D., Gillespie, G. L., Succop, P., & Mentzel, T.K. (2013). Prospective study of violence against ED workers. *American Journal of Emergency Medicine*, 31(1), 197–205. <https://doi.org/10.1016/j.ajem.2012.07.010>
  20. Wolf, L. A., Perhats, C., Delao, A. M., & Clark, P. R. (2017). Workplace aggression as cause and effect: Emergency nurses' experiences of working fatigued. *International Emergency Nursing*, 33, 48–52. <https://doi.org/10.1016/j.ienj.2016.10.006>
  21. Copeland, D., & Henry, M. (2018). The relationship between workplace violence, perceptions of safety, and professional quality of life among ED staff members in a Level 1 trauma centre. *International Emergency Nursing*, 39, 26–32. <https://doi.org/10.1016/j.ienj.2018.01.006>
  22. Nowrouzi-Kia, B., Isidro, R., Chai, E., Usuba, K., & Chen, A. (2018). Antecedent factors in different types of workplace violence against nurses: A systematic review. *Aggression and Violent Behavior*, 44, 1–7. <https://doi.org/10.1016/j.avb.2018.11.002>
  23. Gillespie, G. L., Bresler, S., Gates, D. M., & Succop, P. (2013). Posttraumatic stress symptomatology among emergency department workers following workplace aggression. *Workplace Health & Safety*, 61(6), 247–254. <https://doi.org/10.1177/216507991306100603>
  24. Lanctôt, N., & Guay, S. (2014). The aftermath of workplace violence among healthcare workers: A systematic literature review of the consequences. *Aggression and Violent Behavior*, 19(5), 492–501. <https://doi.org/10.1016/j.avb.2014.07.010>
  25. Gates, D. M., Gillespie, G. L., & Succop, P. (2011). Violence against nurses and its impact on stress and productivity. *Nursing Economics*, 29(2), 59–66. Retrieved from <https://pdfs.semanticscholar.org/0917/>

# Position Statement

930 E. Woodfield Road, Schaumburg, IL 60173 | 800.900.9659 | ena.org

[6fbbd2b4f7a1d2804befe997fdef55d37a53.pdf?\\_ga=2.257857101.733482224.1564579751-1643752325.1564579751](https://doi.org/10.1111/j.1365-2834.2009.00999.x)

26. Spence Laschinger, H. K., Leiter, M., Day, A., & Gilin, D. (2009). Workplace empowerment, incivility, and burnout: Impact on staff nurse recruitment and retention outcomes. *Journal of Nursing Management*, 17(3), 302–311. <https://doi.org/10.1111/j.1365-2834.2009.00999.x>
27. The Joint Commission. (2016). Bullying has no place in health care. *Quick Safety*, 24. Retrieved from [https://www.jointcommission.org/assets/1/23/Quick\\_Safety\\_Issue\\_24\\_June\\_2016.pdf](https://www.jointcommission.org/assets/1/23/Quick_Safety_Issue_24_June_2016.pdf)
28. Oyeleye, O., Hanson, P., O'Connor, N., & Dunn, D. (2013). Relationship of workplace incivility, stress, and burnout on nurses' turnover intentions and psychological empowerment. *JONA: The Journal of Nursing Administration*, 43(10), 536–542. <https://doi.org/10.1097/NNA.0b013e3182a3e8c9>
29. Hogarth, K. M., Beattie, J., & Morphet, J. (2016). Nurses' attitudes towards the reporting of violence in the emergency department. *Australasian Emergency Nursing Journal*, 19(2), 75–81. <https://doi.org/10.1016/j.aenj.2015.03.006>
30. Arnetz, J. E., Hamblin, L., Ager, J., Luborsky, M., Upfal, M. J., Russell, J., & Essenmacher, L. (2015). Underreporting of workplace violence: Comparison of self-report and actual documentation of hospital incidents. *Workplace Health & Safety*, 63(5), 200–210. <https://doi.org/10.1177%2F2165079915574684>
31. Lynette, J., Echevarria, I., Sun, E., & Ryan, J. G. (2016). Incivility across the nursing continuum. *Holistic Nursing Practice*, 30(5), 263–268. <https://doi.org/10.1097/HNP.000000000000167>
32. Wolf, L., Delao, A., & Perhats, C. (2014). Nothing changes, nobody cares: Understanding the experience of emergency nurses physically or verbally assaulted while providing care. *Journal of Emergency Nursing*, 40(4), 305–310. <https://doi.org/10.1016/j.jen.2013.11.006>
33. Wolf, L. A., Perhats, C., Clark, P. R., Moon, M. D., & Zavotsky, K. E. (2018). Workplace bullying in emergency nursing: Development of a grounded theory using situational analysis. *International Emergency Nursing*, 39, 33–39. <https://doi.org/10.1016/j.ienj.2017.09.002>
34. The Joint Commission. (2019, February). Preventing violence in the health care setting. Sentinel Event Alert, (45) 1–3. Retrieved from [https://www.jointcommission.org/assets/1/18/SEA\\_45\\_Preventing\\_Violence\\_in\\_HCOs\\_REV\\_2\\_14\\_19.pdf](https://www.jointcommission.org/assets/1/18/SEA_45_Preventing_Violence_in_HCOs_REV_2_14_19.pdf)
35. Gillespie, G. L., Gates, D. M., Mentzel, T., Al-Natour, A., & Kowalenko, T. (2013). Evaluation of a comprehensive ED violence prevention program. *Journal of Emergency Nursing*, 39(4), 376–383. <https://doi.org/10.1016/j.jen.2012.12.010>
36. Lenaghan, P. A., Cirrincione, N. M., & Henrich, S. (2018). Preventing emergency department violence through design. *Journal of Emergency Nursing*, 44(1), 7–12. <https://doi.org/10.1016/j.jen.2017.06.012>
37. Hyland, S., Watts, J., & Fry, M. (2016). Rates of workplace aggression in the emergency department and nurses' perceptions of this challenging behaviour: A multimethod study. *Australasian Emergency Nursing Journal*, 19(3), 143–148. <https://doi.org/10.1016/j.aenj.2016.05.002>
38. Arnetz, J. E., Hamblin, L., Russell, J., Upfal, M. J., Luborsky, M., Janisse, J., & Essenmacher, L. (2017). Preventing patient-to-worker violence in hospitals: Outcome of a randomized controlled intervention. *Journal of Occupational and Environmental Medicine*, 59(1), 18–27. <https://doi.org/10.1097/JOM.0000000000000909>
39. Occupational Safety and Health Administration. (n.d.). Workplace violence. Retrieved from <https://www.osha.gov/SLTC/workplaceviolence/>
40. Bowie, V. Gill, M. L., & Fisher, B. (2002). *Violence at work: Causes, patterns and prevention*. Devon, OR.



# Position Statement

930 E. Woodfield Road, Schaumburg, IL 60173 | 800.900.9659 | ena.org

Willian.

41. Morphet, J., Griffiths, D., Beattie, J., Reyes, D. V., & Innes, K. (2018). Prevention and management of occupational violence and aggression in healthcare: A scoping review. *Collegian*, 25(6), 621-632.

## Authored by

Lisa Wolf, PhD, RN, CEN, FAEN  
Cydne Perhats, MPH, CHES  
Altair Delao, MPH

## Reviewed by

### 2019 ENA Position Statement Committee

Carla B. Brim, MN, RN, ARNP, CNS, CEN, PHCNS-BC, FAEN  
Judith Carol Gentry, MHA, BSN, RN, CEN, CPEN, CFRN, CTRN, CNML, NE-BC, RN-BC  
Sue L. Leaver, MSN, RN, CEN  
AnnMarie R. Papa, DNP, RN, CEN, NE-BC, FAEN, FAAN  
Matthew Edward Proud, DNP, RN, CEN  
Cheryl Lynn Riwitis, MSN, RN, FNP, EMT-B, CEN, CFRN, FNP-BC, TCRN, FAEN  
Kathryn Starr Rogers, DNP, RN, CEN, CPEN, CPHQ, NEA-BC, TCRN  
Elizabeth L. Stone, PhD, RN, CPEN  
Jennifer Schieferle Uhlenbrock, DNP, MBA, RN, TCRN  
Justin Winger, PhD, MA, BSN, RN, Chairperson  
Mary Ellen Zaleski, DNP, RN, CEN, RN-BC, FAEN

### 2019 ENA Board of Directors Liaison

Gordon Lee Gillespie, PhD, DNP, RN, CEN, CPEN, CNE, PHCNS-BC, FAEN, FAAN

### 2019 ENA Staff Liaison

Monica Escalante Kolbuk, MSN, RN, CEN

Developed: 1991

Approved by the ENA Board of Directors: July 1991  
Revised and Approved by the ENA Board of Directors: September 1993  
Revised and Approved by the ENA Board of Directors: April 1995  
Revised and Approved by the ENA Board of Directors: September 1997  
Revised and Approved by the ENA Board of Directors: December 1999  
Revised and Approved by the ENA Board of Directors: July 2001  
Revised and Approved by the ENA Board of Directors: April 2006  
Revised and Approved by the ENA Board of Directors: September 2008





# Position Statement

930 E. Woodfield Road, Schaumburg, IL 60173 | 800.900.9659 | ena.org

Revised and Approved by the ENA Board of Directors: December 2010  
Revised and Approved by the ENA Board of Directors: October 2014  
Revised and Approved by the ENA Board of Directors: September 2019

This statement updates and replaces, *Violence in the Emergency Setting*, (2014).

©Emergency Nurses Association, 2019.

This position statement, including the information and recommendations set forth herein, reflects ENA's current position with respect to the subject matter discussed herein based on current knowledge at the time of publication. This position statement is only current as of its publication date and is subject to change without notice as new information and advances emerge. The positions, information and recommendations discussed herein are not codified into law or regulations. In addition, variations in practice, which take into account the needs of the individual patient and the resources and limitations unique to the institution, may warrant approaches, treatments and/or procedures that differ from the recommendations outlined in this position statement. Therefore, this position statement should not be construed as dictating an exclusive course of management, treatment or care, nor does adherence to this position statement guarantee a particular outcome. ENA's position statements are never intended to replace a practitioner's best nursing judgment based on the clinical circumstances of a particular patient or patient population. Position statements are published by ENA for educational and informational purposes only, and ENA does not "approve" or "endorse" any specific sources of information referenced herein. ENA assumes no liability for any injury and/or damage to persons or property arising out of or related to the use of or reliance on any position statement.